

## Exercise Problem Sets 1

Sept. 13, 2019

**Problem 1.** Let  $\theta$  be a real number such that  $t = \tan \frac{\theta}{2}$  also be a real number. Show that

$$\sin \theta = \frac{2t}{1+t^2} \quad \text{and} \quad \cos \theta = \frac{1-t^2}{1+t^2}.$$

**Problem 2.** (這題雖不指定上臺講解，但請所有同學都能好好閱讀證明，了解整個流程為什麼就證明了  $\lim_{x \rightarrow c} f(x) = 0$ ) Let  $f : (0, \infty) \rightarrow \mathbb{R}$  be defined by

$$f(x) = \begin{cases} \frac{1}{p} & \text{if } x = \frac{q}{p}, p, q \in \mathbb{N} \text{ and } (p, q) = 1, \\ 0 & \text{if } x \text{ is irrational.} \end{cases}$$

*Proof.* Let  $\varepsilon > 0$  be given. Then there exists a prime number  $p$  such that  $\frac{1}{p} < \varepsilon$ . Let  $q_1, q_2, \dots, q_n$  be rational numbers in  $(\frac{c}{2}, \frac{3c}{2})$  satisfying

$$q_j = \frac{s}{r}, (r, s) = 1, 1 \leq r \leq p,$$

and define  $\delta = \frac{1}{2} \min \left( \{|c - q_1|, |c - q_2|, \dots, |c - q_n|\} \setminus \{0\} \right)$ . Then  $\delta > 0$ . Suppose that  $x$  satisfies that  $0 < |x - c| < \delta$ .

1. If  $x \in \mathbb{Q}^c$ , then  $f(x) = 0$  which shows that  $|f(x)| < \varepsilon$ .
2. If  $x \in \mathbb{Q}$ , then  $x = \frac{s}{r}$  for some natural numbers  $r, s$  satisfying  $(r, s) = 1$ . By the choice of  $\delta$ , we find that  $r > p$ ; thus

$$|f(x)| = \frac{1}{r} < \frac{1}{p} < \varepsilon.$$

In either case,  $|f(x)| < \varepsilon$ ; thus we establish that

$$|f(x) - 0| < \varepsilon \quad \text{whenever} \quad 0 < |x - c| < \delta.$$

Therefore,  $\lim_{x \rightarrow c} f(x) = 0$ . □

**Problem 3.** Let  $f$  be given in Problem 2 and  $g : \mathbb{R} \rightarrow \mathbb{R}$  be defined by

$$g(x) = \begin{cases} f(x) & \text{if } x > 0, \\ f(-x) & \text{if } x < 0, \\ 1 & \text{if } x = 0. \end{cases}$$

Find  $\lim_{x \rightarrow 0} g(x)$ .