

## Exercise Problem Sets 5

Oct. 10. 2019

**Problem 1.** Note that in class we have introduced two new functions “arcsin” and “arccos” whose graphs are (the blue and green) part of the curve consisting of points  $(x, y)$  satisfying  $\sin y = x$  and  $\cos y = x$ , respectively, given below




Figure 1: The graph of functions  $y = \arcsin x$  and  $y = \arccos x$

1. Find the domain and the range of the two functions arcsin and arccos.
2. Show that  $\sin(\arcsin x) = x$  for all  $x$  in the domain of arcsin and  $\cos(\arccos x) = x$  whenever  $x$  in the domain of arccos.
3. Is it true that  $\arcsin(\sin x) = x$  or  $\arccos(\cos x) = x$ ?
4. Find  $\sin(\arccos x)$  and  $\cos(\arcsin x)$ .
5. Show that  $\left. \frac{d}{dx} \right|_{x=c} (\arcsin x + \arccos x) = 0$  for all  $c$  in both domains.
6. Find  $\frac{d}{dx} \arcsin \frac{1}{x}$  and  $\frac{d}{dx} (\arccos x)^2$ .

**Problem 2.** The function arctan is defined similarly to functions arcsin and arccos: consider the collection of all points  $(x, y)$  satisfying  $\tan y = x$  (see the figure below), and the blue part is the graph of a function called “arctan”.




Figure 2: The graph of function  $y = \arctan x$

1. Find the domain and the range of the function  $\arctan$ .
2. Show that  $\tan(\arctan x) = x$  for all  $x$  in the domain of  $\arctan$ .
3. Is it true that  $\arctan(\tan x) = x$  for all  $x$  in the domain of  $\tan$ ?
4. Find  $\frac{d}{dx} \arctan x$ .

**Problem 3.** Find  $\frac{dy}{dx}$  and  $\frac{d^2y}{dx^2}$  if  $\sin(x + y) = y^2 \cos x$ .

**Problem 4.** The line that is normal to the curve  $x^2 + 2xy - 3y^2 = 0$  at  $(1, 1)$  intersects the curve at what other point?

**Problem 5.** Show that the sum of the  $x$ - and  $y$ -intercepts of any tangent line to the curve  $\sqrt{x} + \sqrt{y} = \sqrt{c}$  is equal to  $c$ .

**Problem 6.** The Bessel function of order 0, denoted by  $y = J_0(x)$ , satisfies the differential equation

$$xy'' + y' + xy = 0$$

for all values of  $x$  and its value at 0 is  $J_0(0) = 1$ .

1. Find  $J_0'(0)$ .
2. Use implicit differentiation to find  $J_0''(0)$ .