

## Exercise Problem Sets 2

Sept. 20. 2019

**Problem 1.** Let  $f$  be a function defined on an open interval containing  $c$  (except possibly at  $c$ ).

1. Prove that if  $\lim_{x \rightarrow c} f(x) = L$ , then  $\lim_{x \rightarrow c} |f(x)| = |L|$ .
2. Prove that  $\lim_{x \rightarrow c} f(x) = L$  if and only if  $\lim_{x \rightarrow c} |f(x) - L| = 0$ .

**Problem 2.** Let  $f$  be a function defined on an open interval containing  $c$  and  $\lim_{x \rightarrow c} f(x)$  exists. Show that there exist  $\delta > 0$  and  $M > 0$  such that

$$|f(x)| \leq M \quad \text{whenever} \quad |x - c| < \delta.$$

**Problem 3.** Let  $f, g$  be a function defined on an open interval containing  $c$  (except possibly at  $c$ ), and  $f(x) \leq g(x)$  for all  $x \neq c$ . Prove that if  $\lim_{x \rightarrow c} f(x) = L$  and  $\lim_{x \rightarrow c} g(x) = K$  both exist, then  $L \leq K$ .

**Problem 4.** 1. Suppose that  $\lim_{x \rightarrow 2} \frac{f(x) - 5}{x - 2} = 3$ . Find  $\lim_{x \rightarrow 2} f(x)$ .

2. Suppose that  $\lim_{x \rightarrow 2} \frac{f(x) - 5}{x - 2} = 4$ . Find  $\lim_{x \rightarrow 2} f(x)$ .

3. Suppose that  $\lim_{x \rightarrow c} \frac{f(x) - p(x)}{x - c} = L$  exists, where  $p$  is a polynomial function. Find  $\lim_{x \rightarrow c} f(x)$ .

**Problem 5.** Suppose that you are given  $\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$ . Compute the following limits:

1.  $\lim_{x \rightarrow 0} \frac{\sin(x^2)}{x}$ .      2.  $\lim_{x \rightarrow 0} \frac{x \sin x}{1 - \cos x}$ .      3.  $\lim_{x \rightarrow 0} \frac{\sin(\sin(\sin x))}{x}$ .

4.  $\lim_{x \rightarrow 0} \frac{\sin(x + c) - \sin c}{x}$ , where  $c$  is a real number.

**Problem 6.** Show that  $\lim_{x \rightarrow 0^+} x^{\frac{3}{4}} \cos \frac{1}{x^2} = 0$  using (1)  $\epsilon$ - $\delta$  definition and (2) the Squeeze theorem.

**Problem 7.** 1. Find the limits  $\lim_{x \rightarrow 2^+} \frac{x^2 + x - 6}{|x - 2|}$  and  $\lim_{x \rightarrow 2^-} \frac{x^2 + x - 6}{|x - 2|}$ . Determine whether the limit

$$\lim_{x \rightarrow 2} \frac{x^2 + x - 6}{|x - 2|} \text{ exists or not.}$$

2. Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be defined by

$$f(x) = \begin{cases} a + \sin(x - 2) & \text{if } x > 2, \\ x^2 - 3x + b & \text{if } x \leq 2. \end{cases}$$

Find the relation between  $a$  and  $b$  so that the limit  $\lim_{x \rightarrow 2} f(x)$  exists.

3. Let  $g : \mathbb{R} \rightarrow \mathbb{R}$  be defined by

$$g(x) = \begin{cases} 1 + \sin(x - 2) & \text{if } x > 2, \\ x^2 - 3x + 3 & \text{if } x \leq 2. \end{cases}$$

Find the limit  $\lim_{x \rightarrow 2} \frac{g(x) - 1}{x - 2}$  using the left limit and right limit criteria. You can use the limit in Problem 5.

**Problem 8. True or False:** Determine whether the following statements are true or false. If it is true, prove it. Otherwise, give a counter-example.

1. If  $f$  and  $g$  are functions such that  $\lim_{x \rightarrow c^+} g(x) = K$ ,  $\lim_{y \rightarrow K^+} f(y) = f(K)$ , then

$$\lim_{x \rightarrow c^+} (f \circ g)(x) = f(K).$$

How about if  $x \rightarrow c^+$  and  $y \rightarrow K^+$  are replaced by  $x \rightarrow c^-$  and  $y \rightarrow K^-$ , respectively?

2. Let  $f, g$  be a function defined on an open interval containing  $c$  (except possibly at  $c$ ), and  $f(x) < g(x)$  for all  $x \neq c$ . If  $\lim_{x \rightarrow c} f(x) = L$  and  $\lim_{x \rightarrow c} g(x) = K$  both exist, then  $L < K$ .